

# ON SOME COMBINATORIAL RESULTS OF COLLAPSE IN PARTIAL TRANSFORMATION SEMIGROUPS

<sup>1</sup>Mbah, M. A., <sup>2</sup>Ndubuisi, R. U. and <sup>3</sup>Achaku D. T.

<sup>1,3</sup>Department of Mathematics, Federal University of Lafia P.M.B 146 Lafia

<sup>1,3</sup>Email: <sup>1</sup>mosembah@yahoo.com, <sup>3</sup>achakudt66@gmail.com

<sup>3</sup>Department of Mathematics, Federal University of Technology, Owerri

<sup>2</sup>Email: u\_ndubuisi@yahoo.com

## Abstract.

This paper considers the semigroup of partial transformation  $P_n$  and investigated the elements of collapse given by  $c(\alpha) = |C(\alpha)| = \left| \bigcup_{t \in Im\alpha} \{t\alpha^{-1} \geq 2\} \right|$ . Consequently, the formular for the total number of collapsible elements for  $|t\alpha^{-1}| = 2$  and  $|t\alpha^{-1}| = 3$  in  $P_n$  is presented.

**Keywords:** *Collapse, idempotent, Nilpotent, Partial transformation semigroup.*

## 1 INTRODUCTION

The theory of semigroup is a natural generalization of group theory which has been popular topic in algebra for many years. The most natural occuring semigroups of this generalization is the transformation semigroup, see [3], [5], [7].

A set  $S$  with a binary operation  $\tau$  is called a semigroup if  $\tau$  is associative on  $S$ . In other words, a semigroup is an algebraic system consisting of a non-empty set together with an associative binary operation that is it is obtained from the set of all transformation on a given set. If  $\gamma$  and  $\lambda$  are transformations on  $S$  then, their product  $\gamma\lambda$  is defined as;

$$(x)(\gamma\lambda) = ((x)\gamma)\lambda .$$

Hence we see clearly, that associative law holds for transformation on the set  $S$ .

Let  $X_n = \{1, 2, \dots, n\}$ , then a (partial) transformation  $\alpha : Dom\alpha \subseteq X_n \rightarrow Im\alpha$  is said to be full or total if  $Dom\alpha = X_n$  otherwise it is called strictly partial. The set of all partial transformation on  $n$ -object forms a semigroup under the usual composition of transformations. In [16], Wagner stated that the composition of partial transformation is a special case of the multiplication of binary relations. He recognized also that the domain of composition of two partial transformations may be the empty set so he introduces an empty transformation to take account of this. With the addition of this empty transformation, the composition of partial transformations of a set becomes an

everywhere defined associative binary operation.

The combinatorial properties of  $S_n$  (the partial symmetric semigroup) have been studied over long period and many interesting results have emerged. In particular, the number of permutations of  $X_n$  having exactly  $k$  fixed points and their generating functions are known.

Various enumerative problems of an essentially combinatorial nature have been considered for certain classes of semigroups of transformations. For example, it is well known that  $P_n$  has order  $(n+1)^n$ . Also the number of idempotents and nilpotent for  $P_n$  are given by

$|E(P_n)| = \sum_{r=0}^n \binom{n}{r} (r+1)^{n-r}$  and  $|N(P_n)| = (n+1)^{n-1}$  respectively. The first is usually connected to

Garba [2] and the second can easily be deduced from Laradji and Umar [5].

A study on partial transformation semigroup which we might find interesting in this paper is knowing the total number of collapsible elements  $|c(\alpha)|$ . In [4], Howie defined a collapsible element as  $c(\alpha) = \bigcup_{t \in \text{Im}\alpha} \{t\alpha^{-1} : |t\alpha^{-1}| \geq 2\}$ .

The collapsible element for  $|t\alpha^{-1}| = 2$  and  $|t\alpha^{-1}| = 3$  from  $n \geq 2$  in  $T_n$  was studied by Adeniji and Makanjuola [1]. Other interesting results concerning partial transformation semigroup can be found in [8]. Again for a non-collapsible elements of  $P_n$  that is the partial symmetric semigroup  $S_n$  we have  $(n+1)!$  for  $c(\alpha)$ , this is a case where no image appears more than once.

In this paper we investigate collapse in partial transformation semigroup and obtain formula for the total number of collapsible element for  $|t\alpha^{-1}| = 2$  and  $|t\alpha^{-1}| = 3$  for  $n \geq 2$  in  $P_n$ . Finally, we generalize the formula for  $|t\alpha^{-1}| = 2$  and  $|t\alpha^{-1}| = 3$ .

## 2 PRELIMINARIES

For standard terms and concepts in transformation semigroup theory see [5] or [7]. We now recall notations and definitions that will be used in this paper.

**Definition 2.1.** Consider  $X_n = \{1, 2, \dots, n\}$  and let  $\alpha: X_n \rightarrow X_n$  be a partial transformation. By  $\text{Dom}\alpha$  and  $\text{Im}\alpha$  we mean the domain of the mapping  $\alpha$  and image set or range of the mapping  $\alpha$  respectively.

As in [14] let  $\alpha \in P_n$  be denoted by  $\alpha = \begin{pmatrix} T_1 & T_2 & \dots & T_r \\ t_1 & t_2 & \dots & t_r \end{pmatrix}$ , where  $T_1, T_2, \dots, T_r$  are pairwise disjoint subsets of  $X_n$  called the blocks of  $\alpha$  with  $T_i\alpha = t_i$ . We can be easily seen that the image of  $\alpha$  is  $\text{Im}\alpha = \{t_1, t_2, \dots, t_r\}$  and  $T_1 \cup T_2 \cup \dots \cup T_r = X_n$ . If  $t_i \in T_i$ , we say that  $T_i$  is stationary.

Otherwise it is non-stationary. If we denote by  $F(\alpha)$  the set  $\{T_1 \in X_n : T_1\alpha = T_1\}$  and by  $f(\alpha)$  its cardinal, then the number of stationary blocks of  $\alpha$  is equal to  $f(\alpha)$ . The element  $\alpha$  is idempotent if and only if all the blocks of  $\alpha$  are stationary or if  $f(\alpha) = |Im\alpha|$ .

**Proposition 2.2.** The set  $P_n$  contains  $(n + 1)^n$  elements.

**Proof.** Each element  $\alpha \in P_n$  is uniquely defined by  $\alpha = \begin{pmatrix} 1 & 2 & \dots & \dots & n \\ k_1 & k_2 & \dots & \dots & k_n \end{pmatrix}$  where  $k_i = \alpha(i)$  if  $i \in dom(\alpha)$  and  $k_i = \emptyset$  if  $i \in dom(\alpha)$  again the element  $k_i$  can be independently chosen from the set  $N \cup \{\emptyset\}$ . Hence the product rule implies  $|P_n| = (n + 1)^n$  elements.

**Lemma 2.3.** For all natural numbers  $n$  and  $t$  we have

$$\sum_{k=t}^n \binom{k-1}{t-1} = \binom{n}{t}$$

**Lemma 2.4.** For all natural numbers  $k, n$  and  $t$  we have

$$\sum_{k=0}^n \binom{k+t-2}{k-1} = \binom{n+t-1}{n-1} = \binom{n+t-1}{t}$$

**Lemma 2.5.** Let  $\{P_i\} \in Dom\alpha$  and  $p_i \in Im\alpha$  such that  $|p_i\alpha - 1| \geq 2$ . If  $p_1 \neq p_2 \neq p_3 \neq \dots \neq p_i$  then,  $\varphi(n, 1) = 0$  and  $\varphi(n, 0) = (n + 1)!$  where  $i = 1, 2, 3, \dots, n, n \in \mathbb{N}$ .

### 3 MAIN RESULT

Our result is shown in the triangle of numbers below and we prove a theorem that establishes the total number of collapsible element for  $c(\alpha)$ ,  $|t\alpha^{-1}| = 2$  and  $|t\alpha^{-1}| = 3$  for  $n \geq 2$  in  $P_n$ .

Triangle of numbers  $C(n/q)$

C(n/q)	0	1	2	3	4	5	$\sum C(n; q)$
	2	0					2
	6	0	3				9
	24	0	36	4			64
	120	0	360	80	65		625
	720	0	3600	1200	2250	6	7776

**Theorem 3.1** Let  $P_n$  be a partial transformation semigroup and  $q = C(\alpha)$  defined by  $c(\alpha) = |C(\alpha)| = |\bigcup \{t\alpha^{-1} \geq 2\}|$ , then the following is true:

i) if  $q = 0, C(\alpha) = (n+1)!$  .

ii) if  $q = 2, C(\alpha) = \frac{(n+1)!}{2!} \binom{n}{q}$ .

iii) if  $q = 3, C(\alpha) = \frac{(n+1)!}{3!} \binom{n}{q}$ .

**Proof.** i) When  $q = 0$ , the element in the semigroup of partial transformation with zero element will be the permutation of all the element which is  $n+1$ , adding the zero element. Hence  $c(\alpha) = (n+1)!$

ii) When  $q = 2$ , the number of element will be the number of ways of choosing two element from the  $n$  elements which is  $\binom{n}{2}$  and the arranging of the elements which is  $\frac{n+1}{2}$ . So the total number will be  $\frac{(n+1)!}{2!} \binom{n}{q}$ .

iii) When  $q = 3$  the number of element will be the number of ways of combining three elements from the  $n$  elements which is  $\binom{n}{3}$  and the arranging of the elements which is  $\frac{n+1}{3}$  so that the total number is  $\frac{(n+1)!}{3!} \binom{n}{q}$ .

Hence we give a general form of collapse in partial transformation semigroup for  $n = 0, 2, 3$  as

$$c(\alpha) = \frac{(n+1)!}{q!} \binom{n}{q}.$$

#### 4. CONCLUSION

It has been shown that the number of elements that are collapsible in partial transformation semigroup for values of each of  $q = 2$  and  $q = 3$  can be calculated using  $c(\alpha) = \frac{(n+1)!}{q!} \binom{n}{q}$ .

## REFERENCES

- [1] Adeniji, A.O and Makanjuola, S.O. (2008), Combinatorial Nature of Some Semigroups and collapse in full transformation semigroup. *Afr.j.Comp and ICT*, Vol,No.2,2008.
- [2] Clifford, A.H. and Preston, G. B. (1961). *The algebraic theory of semigroups*, math.surveys No. 7 vol 1 (Amer. Math. Soc., providence, RI).
- [3] Ganyushkin, O and Mazorchuk V. (2009), *Classical Finite transformation Semigroup. An introduction*, Springer.
- [4] Garba, G.U.(1990) Idempotents in partial transformation semigroups *Proc Roy.Soc.Edinburgh* 116A:359-366.
- [5] Higgins, P. N. (1992) *Techniques of Semigroup theory*, Oxford University Press.
- [6] Higgins, P. N. (1993) Combinatorial results for semigroup of order preserving mapping. *Math Proc. Camb.Phil.Soc* (113), 281-296.
- [7] Howie, J.M. (1995), *Fundamentals of Semigroup Theory*, Clarendon Press, Oxford.
- [8] Howie, J.M. (1966) The subsemigroup generated by the idempotents of a full transformation semigroups. *J.London Math Soc* 41:707-716.
- [9] Laradji, A and Umar, A. (2004) "On the number of nilpotents in the Partial symmetric semigroup". *Comm. Algebra* 32:3017-3023.
- [10] Laradji, A and Umar, A. (2004) "On certain finite semigroups of order decreasing transformations I, semigroup forum 69(2) 184-200
- [11] Mbah, M.A.(2014)"Combinatorial results on the Semigroup of Order-Preserving and Order-Decreasing Partial Transformation and Collapse in Partial Transformation", M.Sc Dissertation (Unpublished), Department of Mathematics,University of Ilorin, Nigeria.
- [12] N. V. A. Sloane. *The On-line Encyclopedia of Integer Sequences*, available at <http://www.research.att.com/njas/sequences/>
- [13] Umar, A.(1998), Enumeration of certain finite semigroups of transformations, *Discrete mathematics* 189:291-297.
- [14] Umar, A. (2010) Some Combinatorial Problems in The Theory of Partial Transformation Semigroups. *Algebra and Discrete Math* 9. 115-126.
- [15] Wagner, V.V (1952) "Generalized Groups (Russian)". *Doklady*.