

RECOVERING THE INITIAL CONDITION OF HYPERBOLIC EQUATIONS FROM LATERAL CAUCHY DATA AS A GENERALIZED PROBLEM OF MOMENTS

^{1,2}María B. Pintarelli

¹*Departamento de Matematica de la Facultad de Ciencias Exactas
Universidad Nacional de La Plata, LaPlata -1900. Argentina*

²*Departamento de Ciencias Basicas de la Facultad de Ingenieria
Universidad Nacional de La Plata -1900. Argentina
Corresponding Author Email: mariabpintarelli@gmail.com*

Abstract

It will be shown that to recover the initial condition and finding the solution of a hyperbolic equation with Cauchy conditions can be solved in two steps writing the hyperbolic equation as an integral equation, which can be solved numerically applying the techniques of inverse generalized moments problem. In a first step, the initial condition is found in approximate form, and in a second step we numerically approximate the solution of the hyperbolic equation using the initial condition found.

The method is illustrated with examples.

Keywords: *hyperbolic equation, integral equations, generalized moment problem, inverse problem.*

INTRODUCTION

We want to find $w(x, t)$ and $\varphi(x)$ such that

$$w_{tt} - w_{xx} = f(x, t) \text{ and } w(x, 0) = \varphi(x)$$

with $f(x, t)$ known about a D domain where $D = \{(x, t); a_1 < x < b_1, t > 0\}$. The underlying space is $L^2(D)$ under the conditions

$$w(a_1, t) = f_1(t) \quad w(b_1, t) = f_2(t) \quad t \geq 0$$

$$w_x(a_1, t) = k_1(t) \quad w_x(b_1, t) = k_2(t) \quad t \geq 0$$

$$w_t(x, 0) = g_1(x) \quad a_1 < x < b_1$$

The inverse problems for nonlinear partial differential equations have received considerable attention in the literatura. For example in [1] it is proposed a numerical method to solve an

inverse source problem of computing the initial condition of hyperbolic equations from measurements of Cauchy data. This problem arises in thermo and photo-acoustic tomography in a bounded cavity, in which the reflection of the wave makes the widely-used approaches, such as the time reversal method, not applicable. In order to solve this inverse source problem, we approximate the solution to the hyperbolic equation by its Fourier series with respect to a special orthonormal basis of L^2 .

In [2] is devoted to some inverse boundary problems associated with a time-dependent semilinear hyperbolic equation, where both non linearity and sources (including initial displacement and initial velocity) are unknown. It is shown in several generic scenarios that one can uniquely determine then on linearity and/or the sources by using passive or active boundary observations.

In [3] the main aim of this paper is to solve an inverse source problem for a general nonlinear Hyperbolic equation. Combining the quasi-reversibility method and a suitable Carleman weight function, is defined a map of which fixed point is the solution to the inverse problem.

In [4] it is considered the inverse problem of determining the initial states or the source term of a Hyperbolic equation damped by some non-local time-fractional derivative. This framework is relevant to medical imaging such as thermoacoustic or photoacoustic tomography.

In [5] this paper aims to determine the initial conditions for quasi-linear hyperbolic equations that include non local elements. It is suggested a method where is approaching the solution of the hyperbolic equation by truncating its Fourier series in the time domain with a polynomial-exponential basis.

In [6] this paper, approaches to the numerical recovering of the initial condition in the inverse problem for a nonlinear singularly perturbed reaction–diffusion–advection equation are considered. The feature of the formulation of the inverse problem is the use of additional information about the value of the solution of the equation at the known position of a reaction front, measured experimentally with a delay relative to the initial moment of time. In this case, for the numerical solution of the inverse problem, the gradient method of minimizing the cost functional is applied.

In [7] the inverse problem of restoring the initial condition for the time derivative for the one-dimensional wave equation is considered. As an additional condition, the solution of the wave equation at a finite time is given.

The objective of this work is to show that we can solve the problem using the techniques of inverse moments problem. We focus the study on the numerical approximation.

The interest is not to compare with the existing methods, but to present a different method to my novel criteria, and the one that I have already applied in other cases of partial differential equations under other conditions, for example the Parabolic equation under Cauchy conditions or Dirichlet or Neumann. It turns out that a change in conditions implies a different approach. This is a significant change in the problem statement for its resolution.

The generalized moments problem [8,9,10], is to find a function $f(x)$ about a domain $\Omega \subset \mathbb{R}^d$ that satisfies the sequence of equations

$$\mu_i = \int_{\Omega} g_i(x) f(x) dx \quad i \in \mathbb{N} \quad (1)$$

where N is the set of the natural numbers, $(g_i(x))$ is a given sequence of functions in $L^2(\Omega)$ linearly independent known and the succession of real numbers $\{\mu_i\}_{i \in N}$ are known data.

The moments problem is an ill-conditioned problem in the sense that there may be no solution and if there is no continuous dependence on the given data [8,9,10]. There are several methods to build regularized solutions. One of them is the truncated expansion method [8].

This method is to approximate (1) with the finite moments problem

$$\mu_i = \int_{\Omega} g_i(x) f(x) dx \quad i = 1, 2, \dots, n. \quad (2)$$

where it is considered as approximate solution of to $p_n(x) = \sum_{i=0}^n \lambda_i \phi_i(x)$, and the functions $\{\phi_i(x)\}_{i=1, \dots, n}$ result of orthonormalize $\langle g_1, g_2, \dots, g_n \rangle$ being λ_i the coefficients based on the data μ_i . In the subspace generated by $\langle g_1, g_2, \dots, g_n \rangle$ the solution is stable. If $n \in N$ is chosen in an appropriate way then the solution of (2) it approaches the solution of the problem (1).

In the case where the data μ_i are inaccurate the convergence theorems should be applied and error estimates for the regularized solution (pages 19 - 30 of [8]).

ARTICLE ORGANIZATION

To find $w(x, t)$ and $\varphi(x)$ such that

$$w_{tt} - w_{xx} = f(x, t) \quad \text{and} \quad w(x, 0) = \varphi(x)$$

with $f(x, t)$ known about a D domain where $D = \{(x, t); a_1 < x < b_1, t > 0\}$ under the conditions

$$w(a_1, t) = f_1(t) \quad w(b_1, t) = f_2(t) \quad t \geq 0$$

$$w_x(a_1, t) = k_1(t) \quad w_x(b_1, t) = k_2(t) \quad t \geq 0$$

$$w_t(x, 0) = g_1(x) \quad a_1 < x < b_1$$

we will do it in two steps.

The next section describes the first step.

The section that follows explains the second step.

Then it is explained how the generalized moment problem is solved with the truncated expansion method.

Finally the numerical example and the conclusions.

APPROXIMATION OF THE INITIAL CONDITION $\varphi(x)$ - FIRST STEP

We consider

$$w_{tt} - w_{xx} = f(x, t) \quad (2)$$

the conditions are:

$$w(a_1, t) = f_1(t) \quad w(b_1, t) = f_2(t) \quad t \geq 0$$

$$w_x(a_1, t) = k_1(t) \quad w_x(b_1, t) = k_2(t) \quad t \geq 0$$

$$w_t(x, 0) = g_1(x) \quad a_1 < x < b_1$$

We take as an auxiliary function

$$u(m, r, x, t) = e^{-\frac{m}{b_1}x} e^{-(r+1)t}$$

Note that $u_{xx} = \left(\frac{m}{b_1}\right)^2 u$ and $u_{tt} = (r+1)^2 u$.

We define the vector field

$$F^* = (F_1(w), F_2(w)) = (w_x, -w_t)$$

Since $\text{div}(F^*) = -f(x, t)$ we have to:

$$\iint_D u \text{div}(F^*) dA = \iint_D u (-f(x, t)) dA$$

In addition, as $u \text{div}(F^*) = \text{div}(uF^*) - F^* \cdot \nabla u$, so

$$\iint_D u \text{div}(F^*) dA = \iint_D \text{div}(uF^*) dA - \iint_D F^* \cdot \nabla u dA$$

where $\nabla u = (u_x, u_t)$.

And

$$\iint_D F^* \cdot \nabla u dA = \iint_D (F_1 u_x + F_2 u_t) dA$$

Integrating by parts with respect to x :

$$\begin{aligned} \iint_D F_1 u_x dA &= \int_0^\infty \int_{a_1}^{b_1} F_1 u_x dx dt = \\ &= \int_0^\infty (w(b_1, t) u_x(m, r, b_1, t) - w(a_1, t) u_x(m, r, a_1, t)) dt - \iint_D w u_{xx} dA = \\ &= \int_0^\infty (w(b_1, t) u_x(m, r, b_1, t) - w(a_1, t) u_x(m, r, a_1, t)) dt - \iint_D w \left(\frac{m}{b_1}\right)^2 u dA \end{aligned}$$

Besides

$$\begin{aligned} \iint_D F_2 u_t dA &= \int_0^\infty \int_{a_1}^{b_1} F_2 u_t dx dt = \int_0^\infty \int_{a_1}^{b_1} (-w_t) u_t dx dt = \\ &= - \int_{a_1}^{b_1} w(x, 0) u_t(m, r, x, 0) dx - \int_0^\infty \int_{a_1}^{b_1} w u_{tt} dx dt \end{aligned}$$

then

$$\iint_D F^* \cdot \nabla u dA = \int_0^\infty (w(b_1, t) u_x(m, r, b_1, t) - w(a_1, t) u_x(m, r, a_1, t)) dt -$$

$$\begin{aligned}
& - \iint_D u_{xx} w dA - \int_{a_1}^{b_1} w(x, 0) u_t(m, r, x, 0) dx - \int_0^\infty \int_{a_1}^{b_1} w u_{tt} dx dt = \\
& = \int_0^\infty \left(w(b_1, t) \left(-\frac{m}{b_1}\right) u(m, r, b_1, t) - w(a_1, t) \left(-\frac{m}{b_1}\right) u(m, r, a_1, t) \right) dt - \\
& - \int_{a_1}^{b_1} w(x, 0) (-(r+1)) u(m, r, x, 0) dx - \int_0^\infty \int_{a_1}^{b_1} w u \left(\left(\frac{m}{b_1}\right)^2 - (r+1)^2 \right) dx dt = \\
& = A(m, r) + B(m, r) - \int_0^\infty \int_{a_1}^{b_1} w u \left(\left(\frac{m}{b_1}\right)^2 - (r+1)^2 \right) dx dt
\end{aligned}$$

where

$$\begin{aligned}
A(m, r) &= \int_0^\infty \left(w(b_1, t) \left(-\frac{m}{b_1}\right) u(m, r, b_1, t) - w(a_1, t) \left(-\frac{m}{b_1}\right) u(m, r, a_1, t) \right) dt \\
B(m, r) &= - \int_{a_1}^{b_1} w(x, 0) (-(r+1)) u(m, r, x, 0) dx
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \int_C (u F^*) \cdot n ds = \\
& = \int_{a_1}^{b_1} u(m, r, x, 0) w_t(x, 0) dx - \int_0^\infty u(m, r, b_1, t) w_x(b_1, t) dt - \int_0^\infty u(m, r, a_1, t) w_x(a_1, t) dt = G(m, r) \\
& \therefore \iint u (-f(x, t)) dA = G(m, r) - A(m, r) - B(m, r) + \iint_D w u \left(\left(\frac{m}{b_1}\right)^2 - (r+1)^2 \right) dA
\end{aligned}$$

So if we do $r+1 = \frac{m}{b_1}$:

$$\begin{aligned}
& \int_{a_1}^{b_1} -\left(\frac{m}{b_1}\right) u \left(m, \left(\frac{m}{b_1}\right) - 1, x, 0 \right) w(x, 0) dx = A \left(m, \left(\frac{m}{b_1}\right) - 1 \right) - \\
& - \int_0^\infty u \left(m, \left(\frac{m}{b_1}\right) - 1, b_1, t \right) w_x(b_1, t) dt - \int_0^\infty u \left(m, \left(\frac{m}{b_1}\right) - 1, a_1, t \right) w_x(a_1, t) dt - \\
& - \iint u \left(m, \left(\frac{m}{b_1}\right) - 1, x, t \right) f(x, t) dA = \phi(m) \\
& \therefore \int_{a_1}^{b_1} u \left(m, \left(\frac{m}{b_1}\right) - 1, x, 0 \right) w(x, 0) dx = \frac{\phi(m)}{-\left(\frac{m}{b_1}\right)} = \phi 1(m)
\end{aligned}$$

To solve this integral equation we give integer values to m , $m = 0, 1, 2, \dots, n$

Then

$$\int_{a_1}^{b_1} \varphi(x) H_m(x) dx = \phi_1(m) = \mu_m - - - - - (3)$$

We interpret (3) as a generalized moments problem.

$p_{1n}(x)$ is the numerical approximation found with the truncated expansion method for $\varphi(x) = w(x, 0)$, with $H_m(x) = u\left(m, \left(\frac{m}{b_1}\right) - 1, x, 0\right) = e^{-\frac{m}{b_1}x}$ $m = 0, 1, 2, \dots n$ where n is conveniently chosen.

In section 4 the truncated expansion method will be explained in detail and a theorem will be given that explains what is the accuracy of the approximation found by this method.

APPROACH TO $w(x, t)$ - SECOND STEP

To find an approximation of $w(x, t)$ a similar approach to the previous one is made where $w(x, 0)$ is replaced by $p_{1n}(x)$ and we do not consider $r + 1 = \frac{m}{b_1}$.

We take the auxiliary function $u(m, r, x, t) = e^{-\frac{m}{b_1}x} e^{-(r+1)t}$.

Note that $u_{xx} = \left(\frac{m}{b_1}\right)^2 u$ and $u_{tt} = (r + 1)^2 u$.

We define the vector field $F^* = (F_1(w), F_2(w)) = (w_x, -w_t)$

Since $\text{div}(F^*) = -f(x, t)$ we have to:

$$\iint_D u \text{div}(F^*) dA = \iint_D u (-f(x, t)) dA$$

In addition, as $u \text{div}(F^*) = \text{div}(uF^*) - F^* \cdot \nabla u$, so:

$$\iint_D u \text{div}(F^*) dA = \iint_D \text{div}(uF^*) dA - \iint_D F^* \cdot \nabla u dA$$

Thus

$$\therefore \iint_D u (-f(x, t)) dA = G(m, r) - A(m, r) - B(m, r) + \iint_D w u \left(\left(\frac{m}{b_1} \right)^2 - (r + 1)^2 \right) dA$$

Then

$$\iint_D w u \left(\left(\frac{m}{b_1} \right)^2 - (r + 1)^2 \right) dA = -G(m, r) + A(m, r) + B(m, r) + \iint_D u f(x, t) dA$$

where now

$$B(m, r) = \int_{a_1}^{b_1} p_{1n}(x) (-(r + 1)) u(m, r, x, 0) dx$$

Equivalently

$$\iint_D w(x, t) H_{mr}(x, t) dA = \frac{-G(m, r) + A(m, r) + B(m, r) + \iint_D u f(x, t) dA}{\left(\left(\frac{m}{b_1} \right)^2 - (r + 1)^2 \right)} = \phi(m, r) = \mu_{mr} - - (4)$$

where

$$H_{m,r}(x) = u(m, r, x, t)$$

We can consider (4) as a two-dimensional generalized moment problem if we discretize giving m and r non-negative integer values $m = 0, 1, 2, \dots, n_1$; $r = 0, 2, \dots, n_2$, where n_1 and n_2 are conveniently chosen.

An approximation $p_{2n}(x, t)$ is found by the truncated expansion method for $w(x, t)$ where $n = n_1 \times n_2$.

It could happen that when discretizing the function ϕ we find that it is discontinuous in some integer values. In that case we can apply the following:

If we are in the case of a two-dimensional problem, we take a base:

$$\psi_{ij}(m, r) = m^i r^j e^{-(m+r)} \quad i = 0, 1, \dots, n_1 \quad j = 0, 1, 2, \dots, n_2$$

and we multiply both members of (4) by $\psi_{ij}(m, r)$ and we integrate with respect to m and r .

We have then the generalized moments problem

$$\int_{a_1}^{b_1} \int_0^\infty w(x, t) K_{ij}(x, t) = \mu_{ij} \text{ --- (5)}$$

where

$$\begin{aligned} \mu_{ij} &= \int_{a_1}^{b_1} \int_0^\infty \phi(m, r) \psi_{ij}(m, r) dm dr \\ K_{ij}(x, t) &= \int_{a_1}^{b_1} \int_0^\infty u(m, r, x, t) \psi_{ij}(m, r) dm dr \end{aligned}$$

We apply the truncated expansion method and find a numerical approximation for $w(x, t)$. Analogously if we are in the one-dimensional case. The base would be

$$\psi_i(m) = m^i \quad i = 0, 1, \dots, n$$

The following section details the truncated expansion method for the two-dimensional case. The one-dimensional case is analogous.

SOLUTION OF THE GENERALIZED MOMENTS PROBLEM

We can apply the detailed truncated expansion method in [10] and generalized in [11] and [12] to find an approximation $p_n(x, t)$ for the corresponding finite problem with $i = 0, 1, 2, \dots, n$ where n is the number of moments μ_i . We consider the basis $\phi_i(x, t)$ $i = 0, 1, 2, \dots, n$ obtained by applying the Gram-Schmidt orthonormalization process on $H_i(x, t)$ $i = 0, 1, 2, \dots, n$.

We approximate the solution $w(x, t)$ with [10] and generalized in [11] y [12]:

$$p_n(x, t) = \sum_{i=0}^n \lambda_i \phi_i(x, t)$$

where

$$\lambda_i = \sum_{j=0}^i C_{ij} \mu_j \quad i = 0, 1, 2, \dots, n$$

And the coefficients C_{ij} verify

$$C_{ij} = \left(\sum_{k=j}^{i-1} (-1) \frac{\langle H_i(x, t) | \phi_k(x, t) \rangle}{\|\phi_k(x, t)\|^2} C_{kj} \right) \cdot \|\phi_i(x, t)\|^{-1} \quad 1 < i \leq n; 1 \leq j < i.$$

The terms of the diagonal are $\|\phi_i(x, t)\|^{-1} \quad i = 0, 1, \dots, n$.

The proof of the following theorem is in [12,13].

In [13] the demonstration is made for b_2 finite. If $b_2 = \infty$ instead of taking the Legendre polynomials we take the Laguerre polynomials. En[14] the demonstration is made for the one-dimensional case.

This Theorem gives a measure about the accuracy of the approximation.

Theorem

We consider $b_2 = \infty$.

Sea $\{\mu_i\}_{i=0}^n$ be a set of real numbers and suppose that $f(x, t) \in L^2((a_1, b_1) \times (a_2, b_2))$ for two positive numbers ε and M verify:

$$\sum_{i=0}^n \left| \iint_E H_i(x, t) f(x, t) dt dx - \mu_i \right|^2 \leq \varepsilon^2.$$

$$\iint_E (x f_x^2 + t f_t^2) \text{Exp}[x + t] dt dx \leq M^2 \text{ --- (5).}$$

And it must be fulfilled that

$$t^i f(x, t) \rightarrow 0 \quad \text{si} \quad t \rightarrow \infty \quad \text{para todo} \quad i \in N$$

then

$$\int_{a_1}^{b_1} \int_{a_2}^{\infty} |f(x, t)|^2 dt dx \leq \min_i \left\{ \|C^T C\| \varepsilon^2 + \frac{1}{8(n+1)^2} M^2; i = 0, 1, \dots, n \right\}$$

where C it is a triangular matrix with elements $C_{ij} \quad (1 < i \leq n; 1 \leq j < i)$

and

$$\int_{a_1}^{b_1} \int_{a_2}^{\infty} |f(x, t) - p_n(x, t)|^2 dt dx \leq \|C^T C\| \varepsilon^2 + \frac{1}{8(n+1)^2} M^2.$$

If b_2 it is not infinite then (5) change by

$$\int_{a_2}^{b_2} \int_{a_1}^{b_1} ((b_1 - a_1)^2 f_x^2 + (b_2 - a_2)^2 f_t^2) dx dt \leq M^2$$

NUMERICAL EXAMPLES

Example 1

We consider the equation

$$w_{tt} - w_{xx} = 0 \quad \text{in } 1 < x < 3; \quad t > 0$$

whose solution is: $w(x, t) = e^{-x-3t}$ $\varphi(x) = e^{-x}$

$\phi(m, r)$ and $\phi(m)$ they are continuous.

We take $n=5$ moments and is approaching $\varphi(x) = w(x, 0)$ where the accuracy is

$$\sqrt{\int_1^3 (p_{15}(x) - \varphi(x))^2 dx} = 4.72331 \times 10^{-12}$$

In the Fig.1 the graphics of: $p_{15}(x)$ (magenta color) and $\varphi(x)$ (light blue color) are superimposed.

We take $n = 9$ moments and is approaching $w(x, t)$ where the accuracy is

$$\sqrt{\int_1^3 \int_0^\infty (p_{29}(x) - w(x, t))^2 dt dx} = 0.00488935$$

In the Fig. 2 the graphics of: $p_{29}(x)$ (magenta color) and $\varphi(x)$ (light blue color) are superimposed.




Fig. 1: $p_{15}(x)$ and $\varphi(x)$ example 1




Fig. 2: $w(x, t)$ and $p_{29}(x, t)$ example 1

Example 2

We consider the equation

$$w_{tt} - w_{xx} = \frac{1}{4} e^{-t} (4 + \pi^2) \sin\left(\frac{\pi x}{2}\right) \quad \text{in } 0 < x < 1; \quad t > 0$$

whose solution is: $w(x, t) = \sin\left(\frac{\pi x}{2}\right) e^{-t}$ $\varphi(x) = \sin\left(\frac{\pi x}{2}\right)$

$\phi(m, r)$ we find that it is discontinuous in some integer values.

we take a base:

$$\psi_{ij}(m, r) = m^i r^j e^{-(m+r)} \quad i = 0, 1, 2 \quad j = 0, 1, 2$$

and we multiply both members of (4) by $\psi_{ij}(m, r)$ and we integrate with respect to m and r .

We take $n=5$ moments and is approaching $\varphi(x) = w(x, 0)$ where the accuracy is

$$\sqrt{\int_0^1 (p_{15}(x) - \varphi(x))^2 dx} = 0.000975088$$

In the Fig.3 the graphics of: $p_{15}(x)$ (magenta color) and $\varphi(x)$ (light blue color) are superimposed.

$\phi(m, r)$ is continuous. We take $n = 9$ moments and is approaching $w(x, t)$ where the accuracy is

$$\sqrt{\int_0^1 \int_0^\infty (p_{29}(x) - w(x, t))^2 dt dx} = 0.0214964$$

In the Fig. 4 the graphics of: $p_{29}(x)$ (magenta color) and $\varphi(x)$ (light blue color) are superimposed.




Fig. 3: $p_{15}(x)$ and $\varphi(x)$ example 2




Fig. 4: $w(x, t)$ and $p_{29}(x, t)$ example 2

CONCLUSION

A parabolic equation of the form $w_{tt} - w_{xx} = f(x, t)$ where the unknown function $w(x, t)$ is defined in $D = (a_1, b_1) \times (0, \infty)$ under Cauchy conditions with unknown initial condition $\varphi(x) = w(x, 0)$ it can be solved numerically by applying inverse moment problem techniques in two steps.

1. first the partial derivatives equation is written as an integral equation

$$\begin{aligned} \int_{a_1}^{b_1} u\left(m, \frac{m}{b_1} - 1, x, 0\right) w(x, 0) dx &= A\left(m, \frac{m}{b_1} - 1\right) - \\ - \int_0^\infty u\left(m, \frac{m}{b_1} - 1, b_1, t\right) w_x(b_1, t) dt &- \int_0^\infty u\left(m, \frac{m}{b_1} - 1, a_1, t\right) w_x(a_1, t) dt - \\ - \iint u\left(m, \frac{m}{b_1} - 1, x, t\right) f(x, t) dA &= \frac{\phi(m)}{-\left(\frac{m}{b_1}\right)} = \phi 1(m) \end{aligned}$$

To solve this integral equation we give integer values to m : $m = 0, 1, 2, \dots, n$

Then $\int_{a_1}^{b_1} \varphi(x) H_m(x) dx = \phi(m) = \mu_m$. We interpret as a generalized moments problem, $p_{1n}(x)$ is the numerical approximation with the truncated expansion method for $\varphi(x) = w(x, 0)$, with

$H_m(x) = u\left(m, \frac{m}{b_1} - 1, x, 0\right) = e^{-\frac{m}{b_1}x}$ $m = 0, 1, 2, \dots, n$, where n is conveniently chosen.

2. To find an approximation of $w(x, t)$ we consider:

$$\iint_D w(x, t) H_{mr}(x, t) dA = \frac{-G(m, r) + A(m, r) + B(m, r) + \iint_D u f(x, t) dA}{\left(\left(\frac{m}{b_1}\right)^2 - (r + 1)^2\right)} = \phi(m, r) = \mu_{mr}$$

where $H_{m,r}(x) = u(m, r, x, t)$. We can consider it as a two-dimensional generalized moment problem if we discretize giving m and r non-negative integer values $m = 0, 1, 2, \dots, n_1$; $r = 0, 2, \dots, n_2$, where n_1 and n_2 are conveniently chosen. An approximation $p_{2n}(x, t)$ is found by the truncated expansion method for $w(x, t)$ where $n = n_1 \times n_2$.

REFERENCES:

1. Thuy T. Le, Loc H. Nguyen, Thi-Phong Nguyen and William Powell: "The quasi-reversibility method to numerically solve an inverse source problem for hyperbolic equations": arXiv:2011.04855v3 [math.NA] (2021).
2. Yi-Hsuan Lin, Hongyuli, and Xu Liu: "Determining a nonlinear hyperbolic system with unknown sources and nonlinearity": arXiv:2107.10219v2 [math.AP] (2023).
3. Loc H. Nguyen and Michael V. Klibanov: "Carleman estimates and the contraction principle for an inverse source problem for nonlinear hyperbolic equations": arXiv:2108.03500v4 [math.AP] (2022).
4. Xinchu Huang, Yavar Kian, Eric Soccorsi and Masahiro Yamamoto: "Determination of source and initial values for acoustic equations with a time-fractional attenuation": arXiv:2111.05240v1 [math.AP] (2021).
5. Trong D. Dang, Loc H. Nguyen and Huong T. T. Vu: "Determining initial conditions for nonlinear hyperbolic equations with time dimensional reduction and the Carleman contraction principle": arXiv:2312.01179v3 [math.NA] (2024).
6. Dmitry Lukyanenko, Tatyana Yeleskina, Igor Prigorniy, Temur Isaev, Andrey Borzunov and Maxim Shishlenin: "Inverse Problem of Recovering the Initial Condition for a Nonlinear Equation of the Reaction–Diffusion–Advection Type by Data Given on the Position of a Reaction Front with a

Time Delay”: Mathematics, **9(4)**, 342, 12 pages, (2021)

7. Khanlar M. Gamzaev: “Numerical method for restoring the initial condition for the wave equation”: Vestnik Tomskogo gosudarstvennogo universiteta. Matematika i mekhanika – Tomsk State University Journal of Mathematics and Mechanics. 88. pp. 5–13, (2024)

8. D.D. Ang, R. Gorenflo, V.K. Le and D.D. Trong: “Moment theory and some inverse problems in potential theory and heat conduction”: Lectures Notes in Mathematics, Springer-Verlag, Berlin (2002)

9. J.A. Shohat and J.D. Tamarkin: “The problem of Moments”: Mathematic Surveys, Am. Math. Soc., Providence, RI (1943)

10. G. Talenti: “Recovering a function from a finite number of moments”: Inverse Problems: **3**, 501- 517, (1987)

11. M. B. Pintarelli and F. Vericat: “Stability theorem and inversion algorithm for a generalized moment problem”: Far East Journal of Mathematical Sciences: **30**, 253-274, (2008)

12. M.B. Pintarelli and F. Vericat: “Bi-dimensional inverse moment problems”: Far East Journal of Mathematical Sciences: **54**, 1-23, (2011)

13. M.B. Pintarelli: “ Linear partial differential equations of first order as bi-dimensional inverse moment problem”: Applied Mathematics: **6(6)**, 979-989, (2015)

14. M.B. Pintarelli: “ Parabolic partial differential equations as inverse moments problem”: Applied Mathematics: **7 (1)**, 77-99 , (2016)